



Conference Abstract

Basal and $\text{PGF}_{2\alpha}$ -stimulated secretion of pro-inflammatory cytokines from 3T3-L1 adipocyte-like cells

Bilyana Ilieva[‡], Elena Marinova[§], Hristo Gagov[|], Rossitza Konakchieva[‡]

[‡] 1 Sofia University St. Kliment Ohridski, Faculty of Biology, Sofia, Bulgaria

[§] MC ReproBioMed Ltd, Sofia, Bulgaria

[|] Sofia University St. Kliment Ohridski, Faculty of Biology, Sofia, Bulgaria

Corresponding author: Bilyana Ilieva (bmilieva@gmail.com)

Received: 21 Jul 2017 | Published: 01 Aug 2017

Citation: Ilieva B, Marinova E, Gagov H, Konakchieva R (2017) Basal and $\text{PGF}_{2\alpha}$ -stimulated secretion of pro-inflammatory cytokines from 3T3-L1 adipocyte-like cells. BioDiscovery 20: e17757.

<https://doi.org/10.3897/biodiscovery.20.e17757>

Abstract

Adipocytes were recently identified as an important source of endocrine and paracrine mediators, regulating the metabolism and activity of various cell types and body functions. 3T3-L1 preadipocytes are useful model for physiological, pharmacological and cell signaling studies. Differentiation of 3T3-L1 murine fibroblasts into adipocyte-like cells was conducted in presence of IBMX, dexamethasone and insulin and demonstrated by Oil Red O staining of accumulated lipid droplets. Using Inflammatory Multi- Analyte Cytokines ELISArray Kit we investigated the release of cytokines under basal conditions, after $\text{PGF}_{2\alpha}$ treatment for 24 hours to induce pro-inflammatory phenotype, and after $\text{PGF}_{2\alpha}$ treatment and incubation in the presence of L-C-Propargylglycine (PGG, 1 mmol/l), a selective inhibitor of cystathionine-gamma-lyase (CSE). The last combination was used to explore the role of H_2S , released from CSE, for cytokine and H_2O_2 release. We found that $\text{PGF}_{2\alpha}$ strongly increased TNF_{α} secretion from differentiated adipocytes, the latter effect being antagonized by PGG. The CSE inhibitor enhanced IL-6 production and suppressed IL-10 secretion. PGG enhanced H_2O_2 production of in $\text{PGF}_{2\alpha}$ -treated cells. It is concluded that pro-inflammatory phenotype of differentiated 3T3-L1 adipocyte-like cells, induced by PGF

2α is characterized by enhanced TNF_α production which critically depends on the ability of CSE to produce H_2S .

Keywords

adipocytes, paracrine mediators, inflammation, hydrogen sulfide, cystathionine-gamma-lyase (CSE)

Presenting author

Bilyana Ilieva

Presented at

World BioDiscovery Congress 2017

Acknowledgements

This work was supported by SU Scientific Research Fund, project №: 80-10-17/12.04.2017

Conflicts of interest

No